

READ CARTON BEFORE USING THIS PRODUCT

DIRECTIONS FOR USE:

Stir well before use. Apply a thin layer of PROUD-AID across the wound surface using the stirrer – applicator provided. Repeat treatment every 1-3 days until healing is complete. It is recommended that wounds be covered with non-stick gauze and a bandage.

MEAT WITHHOLDING PERIOD: NIL



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

PROUD-AID

WOUND CARE TREATMENT FOR HORSES

100g

ACTIVE CONSTITUENTS

344g/kg Calcium Hydroxide, 29g/kg Copper Sulphate
(as Copper Sulphate Pentahydrate)

TO AID IN THE TREATMENT OF CUTS, WOUNDS AND ABRASIONS AND AID IN THE
TREATMENT AND PREVENTION OF EXCESSIVE GRANULATION TISSUE (**PROUD FLESH**)

STORAGE:

Store below 30°C (Room Temperature).
Keep container tightly closed.

APVMA Approval No. 62714/100g/1009

Batch No:

Expiry:

**pharmachem**
CARING FOR YOUR CREATURE'S COMFORTS

Unit 6, 70 Fison Ave West,
Eagle Farm QLD 4009
phone: (07) 3868 0333