

FOR ANIMAL TREATMENT ONLY

QUIT-ITCH

ANTISEPTIC LOTION

• Bactericide • Fungicide

Active Constituents:

7.5g/L available IODINE (I) (present as 15g/L POVIDONE IODINE and 40g/L IODOPHOR)

BROAD SPECTRUM ANTISEPTIC SKIN TREATMENT FOR SKIN CARE OF HORSES AND DOGS



250mL

An aid in the treatment of ECZEMA, non-specific DERMATOSES and FUNGAL INFECTIONS including RINGWORM (Microsporium spp and Trichophyton spp)

AN EMERGENCY TREATMENT for LACERATIONS, ABRASIONS, WIRE CUTS, BURNS, and other superficial animal wounds

FOR HORSES as an aid in the treatment of QUEENSLAND ITCH, SADDLE ACNE, GIRTH ITCH and GREASY HEEL

DIRECTIONS FOR USE:

CONTRAINDICATIONS: This product is contraindicated in Kittens under 12 weeks old.

WOUNDS: As an emergency treatment for LACERATIONS, ABRASIONS, WIRE CUTS AND BURNS. Dab the wound with cotton wool or gauze soaked with undiluted QUIT-ITCH. Repeat daily for 3-4 days.

SKIN DISEASES: Clean affected area with a solution of 10mL QUIT-ITCH to 200mL warm water. Towel dry and apply undiluted QUIT-ITCH liberally to the affected areas. Repeat daily for 3-4 days.

ALL OVER SHAMPOO WITH QUIT-ITCH: FUNGAL RINGWORM and ECZEMA spread rapidly and in both instances animals should be thoroughly washed all over to prevent spread of infection. Use a solution of 10mL QUIT-ITCH to 200mL warm water. Rub well into the coat and allow the foam to remain for 5 minutes. Rinse, dry and apply undiluted QUIT-ITCH to infected and surrounding areas. HORSES that have been in contact with, or have used the same equipment as an infected horse should also be thoroughly shampooed with diluted QUIT-ITCH daily until the infection is cleared from the stable. This also applies to DOGS kept in kennels.

GREASY HEEL IN HORSES: Most importantly, remove scabs that are indicative of infected heels. Use a solution of 10mL QUIT-ITCH to 200mL of warm water. Rub vigorously to soften and remove scabs. Thoroughly dry and rub undiluted QUIT-ITCH into affected areas twice daily until heels are completely free of infection.

MEAT WITHHOLDING PERIOD: Zero (0) days

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre:
Australia Phone: 13 11 26
New Zealand Phone: 0800 764 766

DISPOSAL: Dispose of container by wrapping with paper and putting in the garbage.

STORAGE: Store below 30°C (Room Temperature)
Keep out of direct sunlight.

APVMA 45430/126271

Distributed by:



Unit 6, 70 Fison Ave West, Eagle Farm QLD 4009
www.pharmachem.com.au



BATCH No:
EXPIRY DATE